

Ibuprofen Dosing Chart

Weight	Age	Concentrated Infant Drops 50 mg/1.25 mL	Children's Suspension 100 mg/5 mL	Junior Strength Chewable 100 mg Each
12-17 lbs	6-11 Months	1.25 mL (50 mg)		
18-23 lbs	12-23 Months	1.875 mL (75 mg)		
24-35 lbs	2-3 Years		1 teaspoon (5 mL)	
36-47 lbs	4-5 Years		1 1/2 teaspoons (7.5 mL)	
48-59 lbs	6-8 Years		2 teaspoons (10 mL)	2 tablets (200 mg)
60-71 lbs	9-10 Years		2 1/2 teaspoons (12.5 mL)	2 1/2 tablets (250 mg)
72-95 lbs	11 Years		3 teaspoons (15 mL)	3 tablets (300 mg)
96+ lbs	12+ Years			4 tablets (400 mg)

AVOID USE IN INFANTS UNDER 6 MONTHS OF AGE

Use only the measuring dropper, syringe or cup that is provided with the medication. One teaspoon = 5 milliliters (5 mL). Do not use a household teaspoon as they may vary in size.

**Ibuprofen Doses May Be Repeated Every 6-8 Hours as Needed.
DO NOT EXCEED 4 DOSES IN 24 HOURS.**

DO NOT USE ASPIRIN TO TREAT FEVER IN CHILDREN.

Dosage information is based on 10mg/kg per ibuprofen dose as recommended by The John Hopkins Hospital: The Harriet Lane Handbook, 16th Edition.

